


# Xavier Pennec


Univ. Côte d'Azur and Inria, France



[http://www-sop.inria.fr/asclepios/cours/Peyresq\\_2019/](http://www-sop.inria.fr/asclepios/cours/Peyresq_2019/)

## Geometric Statistics

*Mathematical foundations  
and applications in  
computational anatomy*



Freely adapted from “Women teaching geometry”, in  
Adelard of Bath translation of Euclid’s elements, 1310.

## 5/ Advanced Stats: empirical estimation and generalized PCA

Ecole d’été de Peyresq, Jul 1-5 2019

*Epione*  
e-patient / e-medicine

UNIVERSITÉ  
CÔTE D’AZUR

*Inria*

# ***Geometric Statistics: Mathematical foundations and applications in computational anatomy***

**Intrinsic Statistics on Riemannian Manifolds**

**Manifold-Valued Image Processing**

**Metric and Affine Geometric Settings for Lie Groups**

**Parallel Transport to Analyze Longitudinal Deformations**

**Advances Statistics: CLT & PCA**

# ***Geometric Statistics: Mathematical foundations and applications in computational anatomy***

Intrinsic Statistics on Riemannian Manifolds

Manifold-Valued Image Processing

Metric and Affine Geometric Settings for Lie Groups

Parallel Transport to Analyze Longitudinal Deformations

## **Advances Statistics: CLT & PCA**

- Estimation of the empirical Fréchet mean & CLT
- Principal component analysis in manifolds
- Natural subspaces in manifolds: barycentric subspaces
- Rephrasing PCA with flags of subspaces

# Several definitions of the mean

## Tensor moments of a random point on M

- $\mathfrak{M}_1(x) = \int_M \overrightarrow{xz} dP(z)$  Tangent mean: (0,1) tensor field
- $\mathfrak{M}_2(x) = \int_M \overrightarrow{xz} \otimes \overrightarrow{xz} dP(z)$  Covariance: (0,2) tensor field
- $\mathfrak{M}_k(x) = \int_M \overrightarrow{xz} \otimes \overrightarrow{xz} \otimes \cdots \otimes \overrightarrow{xz} dP(z)$  k-contravariant tensor field
- $\sigma^2(x) = \text{Tr}_g(\mathfrak{M}_2(x)) = \int_M \text{dist}^2(x, z) dP(z)$  **Variance function**


## Mean value = optimum of the variance

- **Frechet mean** [1944] = (global) minima of p-variance (includes median)
- **Karcher mean** [1977] = local minima
- **Exponential barycenters** = critical points ( $P(C) = 0$ )

$$\mathfrak{M}_1(\bar{x}) = \int_M \overrightarrow{xz} dP(z) = 0 \quad (\text{implicit definition})$$

## Covariance at the mean

$$\mathfrak{M}_2(\bar{x}) = \int_M \overrightarrow{xz} \otimes \overrightarrow{xz} dP(z)$$



# Asymptotic behavior of the mean

## Uniqueness of p-means with convex support

[Karcher 77 / Buser & Karcher 1981 / Kendall 90 / Afsari 10 / Le 11]

- Non-positively curved metric spaces (Aleksandrov): OK [Gromov, Sturm]
- Positive curvature: [Karcher 77 & Kendall 89] concentration conditions:  
Support in a regular geodesic ball of radius  $r < r^* = \frac{1}{2} \min(\text{inj}(M), \pi/\sqrt{\kappa})$

## Bhattacharya-Patangenaru CLT [BP 2005, B&B 2008]

- Under suitable concentration conditions, for IID n-samples:
  - $\bar{x}_n \rightarrow \bar{x}$  (*consistency of empirical mean*)
  - $\sqrt{n} \log_{\bar{x}}(\bar{x}_n) \rightarrow N(0, \bar{H}^{-1} \Sigma \bar{H}^{-1})$  if  $\bar{H} = \int_M \text{Hess}_{\bar{x}}(d^2(y, \bar{x})) \mu(dy)$  invertible

## Questions

- Intelligible expression of Hessian?
- What happens for a small sample size (non-asymptotic behavior)?
- Can we extend results to affine connection spaces?



## Concentration assumptions

- Uniqueness of the mean, support of diameter  $< \varepsilon$

### Riemannian manifold: Karcher & Kendall Concentr. Cond.

- $\text{Supp}(\mu) \subset B(x, r)$  with  $r < \frac{1}{2} \text{inj}(x)$
- $\sup_{x \in B(x, r)} \kappa(x) < \pi^2 / (4r)^2$

### Affine connection spaces: Arnaudon & Li convexity cond.

- $\rho: M \times M \rightarrow R^+$  separating function
  - Separability:  $\rho(x, y) = 0 \Leftrightarrow x = y$
  - Convexity along geodesic:  $\rho(\gamma_1(t), \gamma_2(t)): R \rightarrow R^+$  convex
- p-convex geometry:  $c \text{dist}^p(x, y) \leq \rho(x, y) \leq C \text{dist}^p(x, y)$
- Uniqueness of exponential barycenter (compact support)

# Taylor expansion in manifolds


## The mean is an exponential barycenter

- Tangent mean field:  $\mathfrak{M}_1(x) = \int_M \log_x(z) \mu(dz)$   
has a zero at  $\bar{x}$ . Problem: vector field
- Recentered man field is a mapping of vector spaces

$$N_x(v) = \Pi_{x_v}^x \mathfrak{M}_1(\exp_x(v)) = \int_M \Pi_{x_v}^x \log_{x_v}(y) \mu(dy)$$

has a zero at  $\bar{v} = \log_x(\bar{x})$

## Neighboring log expansion (derived from Gavrilov)



$$\begin{aligned} l_x(v, w) &= w - v + \frac{1}{6} R(w, v)(v - 2w) + \frac{1}{24} \nabla_v R(w, v)(2v - 3w) \\ &\quad + \frac{1}{24} \nabla_w R(w, v)(v - 2w) + O(5) \end{aligned}$$

# Non-Asymptotic behavior of empirical means

## Moments of the Fréchet mean of a n-sample

- Taylor expansions based on [Gavrilov 2007]

- **Unexpected bias** in  $1/n$  on empirical mean (**gradient of curvature-cov.**)

$$\text{bias}(\bar{x}_n) = E(\log_{\bar{x}}(\bar{x}_n)) = \frac{1}{6n} (\mathfrak{M}_2 : \nabla R : \mathfrak{M}_2) + O(\epsilon^5, 1/n^2)$$

- **Concentration rate** modulated by the **curvature-covariance**:

$$\text{Cov}(\bar{x}_n) = E(\log_{\bar{x}}(\bar{x}_n) \otimes \log_{\bar{x}}(\bar{x}_n)) = \frac{1}{n} \mathfrak{M}_2 + \frac{1}{3n} \mathfrak{M}_2 : R : \mathfrak{M}_2 + O(\epsilon^5, 1/n^2)$$

- **Asymptotically infinitely fast CV** for negative curvature
- **No convergence (LLN fails)** at the limit of KKC condition

[XP, Curvature effects on the empirical mean in Manifolds 2019, arXiv:1906.07418 ]



# Constant curvature spaces

- Symmetric spaces: no bias
- Variance is modulated w.r.t. Euclidean:  $Var(\bar{x}_n) = \alpha \frac{\sigma^2}{n}$

## High concentration expansion

- $\alpha = 1 + \frac{2}{3} \left(1 - \frac{1}{d}\right) \left(1 - \frac{1}{n}\right) \kappa \sigma^2 + O(\epsilon^5)$


## Asymptotic CLT expansion





- $\alpha = \left(\frac{1}{d} + \left(1 - \frac{1}{d}\right) \bar{h}\right)^{-2} + O(n^{-2})$

## Archetypal modulation factor

- Uniform distrib on  $S(\bar{x}, \theta) \subset M$   
large  $n$ , large  $d$

- $\alpha = \frac{\tan^2(\sqrt{\kappa} \theta^2)}{\kappa \theta^2}$





# Conclusions

**High concentration expansion very accurate for low theta**

**Asymptotic expansion very accurate for  $n > 10$**

**Main variable controlling the modulation is variance-curvature tensor**

$$R(\mathbf{u}, \mathbf{v}) \mathbf{u} : \mathfrak{M}_2$$

**Main variable controlling the bias**

$$\mathfrak{M}_2 : \nabla \circ R(\mathbf{v}, \mathbf{u}) \mathbf{u} : \mathfrak{M}_2$$

# ***Geometric Statistics: Mathematical foundations and applications in computational anatomy***

Intrinsic Statistics on Riemannian Manifolds

Manifold-Valued Image Processing


Metric and Affine Geometric Settings for Lie Groups

Parallel Transport to Analyze Longitudinal Deformations

## **Advances Statistics: CLT & PCA**


- Estimation of the empirical Fréchet mean & CLT
- **Principal component analysis in manifolds**
- Natural subspaces in manifolds: barycentric subspaces
- Rephrasing PCA with flags of subspaces

# Low dimensional subspace approximation?



Manifold of cerebral ventricles

Etyngier, Keriven, Segonne 2007.



Manifold of brain images

S. Gerber et al, Medical Image analysis, 2009.

- Beyond the 0-dim mean  $\rightarrow$  higher dimensional subspaces
- When embedding structure is already manifold (e.g. Riemannian):  
Not manifold learning (LLE, Isomap,...) but **submanifold learning**
- **Natural subspaces for extending PCA to manifolds?**

# *Tangent PCA (tPCA)*

## Maximize the squared distance to the mean (explained variance)


- Algorithm
  - Unfold data on tangent space at the mean
  - Diagonalize covariance at the mean  $\Sigma(x) \propto \sum_i \overrightarrow{\bar{x}x_i} \overrightarrow{\bar{x}x_i}^t$
- Generative model:
  - Gaussian (large variance) in the horizontal subspace
  - Gaussian (small variance) in the vertical space
- Find the subspace of  $T_x M$  that best explains the variance





# Problems of tPCA

## Analysis is done relative to the mean

- What if the mean is a poor description of the data?
  - Multimodal distributions
  - Uniform distribution on subspaces
  - Large variance w.r.t curvature



Bimodal distribution on  $S^2$



Images courtesy of S. Sommer

# Principal Geodesic / Geodesic Principal Component Analysis

**Minimize the squared Riemannian distance to a low dimensional subspace (unexplained variance)**

- **Geodesic Subspace:**  $GS(x, w_1, \dots, w_k) = \{\exp_x(\sum_i \alpha_i w_i) \text{ for } \alpha \in \mathbb{R}^k\}$ 
  - Parametric subspace spanned by geodesic rays from point  $x$
  - **Beware: GS have to be restricted to be well posed [XP, AoS 2018]**
    - PGA (Fletcher et al., 2004, Sommer 2014)
    - Geodesic PCA (GPCA, Huckeman et al., 2010)
- **Generative model:**
  - Unknown (uniform ?) distribution within the subspace
  - Gaussian distribution in the vertical space

**Asymmetry w.r.t. the base point in  $GS(x, w_1, \dots, w_k)$**

- Totally geodesic at  $x$  only


# Patching the Problems of tPCA / PGA

## Improve the flexibility of the geodesics

- 1D regression with higher order splines [Gu, Machado, Leite, Vialard, Singh, Niethammer, Absil,...]
- Control of dimensionality for n-D Polynomials on manifolds?

## Iterated Frame Bundle Development [HCA, Sommer GSI 2013]

- Iterated construction of subspaces
- Parallel transport in frame bundle
- Intrinsic asymmetry between components



Courtesy of S. Sommer

## Nested “algebraic” subspaces

- Principal nested spheres [Jung, Dryden, Marron 2012]
- Quotient of Lie group action [Huckemann, Hotz, Munk, 2010]
- No general semi-direct product space structure in general Riemannian manifolds

# ***Geometric Statistics: Mathematical foundations and applications in computational anatomy***

Intrinsic Statistics on Riemannian Manifolds

Manifold-Valued Image Processing

Metric and Affine Geometric Settings for Lie Groups

Parallel Transport to Analyze Longitudinal Deformations

## **Advances Statistics: CLT & PCA**

- Estimation of the empirical Fréchet mean & CLT
- Principal component analysis in manifolds
- Natural subspaces in manifolds: barycentric subspaces
- Rephrasing PCA with flags of subspaces


# Affine span in Euclidean spaces

## Affine span of $(k+1)$ points: weighted barycentric equation

$$\begin{aligned}\text{Aff}(x_0, x_1, \dots, x_k) &= \{x = \sum_i \lambda_i x_i \text{ with } \sum_i \lambda_i = 1\} \\ &= \{x \in \mathbb{R}^n \text{ s.t. } \sum_i \lambda_i (x_i - x) = 0, \lambda \in P_k^*\}\end{aligned}$$

### Key ideas:

- ~~□ tPCA, PGA: Look at data points from the mean (mean has to be unique)~~
- Triangulate from several reference:  
**locus of weighted means**



A. Manesson-Mallet. La géométrie Pratique, 1702

# *Barycentric subspaces and Affine span in Riemannian manifolds*

## **Fréchet / Karcher barycentric subspaces (KBS / FBS)**

- Normalized weighted variance:  $\sigma^2(x, \lambda) = \sum \lambda_i \text{dist}^2(x, x_i) / \sum \lambda_i$
- Set of absolute / local minima of the  $\lambda$ -variance
- Works in stratified spaces (may go accross different strata)
  - Non-negative weights: Locus of Fréchet Mean [Weyenberg, Nye]

## **Exponential barycentric subspace and affine span**

- Weighted exponential barycenters:  $\mathfrak{M}_1(x, \lambda) = \sum_i \lambda_i \overrightarrow{xx_i} = 0$
- $\text{EBS}(x_0, \dots, x_k) = \{x \in M^*(x_0, \dots, x_k) \mid \mathfrak{M}_1(x, \lambda) = 0\}$
- Affine span = closure of EBS in  $M$      $\text{Aff}(x_0, \dots, x_k) = \overline{\text{EBS}(x_0, \dots, x_k)}$

## **Questions**

- Local structure: local manifold? dimension? stratification?
- Relationship between  $\text{KBS} \subset \text{FBS}$ ,  $\text{EBS}$  and affine span?

[X.P. Barycentric Subspace Analysis on Manifolds. Annals of statistics. 2018. To appear. arXiv:1607.02833]



# Analysis of Barycentric Subspaces

## Assumptions:

- Restrict to the **punctured manifold**  $M^*(x_0, \dots, x_k) = M \setminus \cup C(x_i)$ 
  - $\text{dist}^2(x, x_i), \log_x(x_i)$  are smooth but  $M^*$  may be split in pieces
- Affinely independent points:  
 $\{\overrightarrow{x_i x_j}\}_{0 \leq i \neq j \leq k}$  exist and are linearly independent for all  $i$

## Local well posedness for the barycentric simplex:

- EBS / KBS are well defined in a neighborhood of reference points
- For reference points in a sufficiently small ball and positive weights:  
unique Frechet = Karcher = Exp Barycenter in that ball: smooth graph of a  $k$ -dim function [proof using Buser & Karcher 81]

## SVD characterization of EBS: $\mathfrak{M}_1(x, \lambda) = Z(x)\lambda = 0$

- SVD:  $Z(x) = [\overrightarrow{xx_0}, \dots, \overrightarrow{xx_k}] = U(x)S(x)V^t(x)$ 
  - $EBS(x_0, \dots, x_k) = \text{Zero level-set of } l > 0 \text{ singular values of } Z(x)$
  - Stratification on the number of vanishing singular values

[X.P. Barycentric Subspace Analysis on Manifolds. Annals of statistics. 2018. To appear. arXiv:1607.02833]

# Analysis of Barycentric Subspaces

**Exp. barycenters are critical points of  $\lambda$ -variance on  $M^*$**

$$\square \nabla \sigma^2(x, \lambda) = -2\mathfrak{M}_1(x, \lambda) = 0$$

$$KBS \cap M^* \subset EBS$$

**Caractérisation of local minima: Hessian (if non degenerate)**

$$H(x, \lambda) = -2 \sum_i \lambda_i D_x \log_x(x_i) = \text{Id} - \frac{1}{3} \text{Ric}(\mathfrak{M}_2(x, \lambda)) + \text{HOT}$$

**Regular and positive pts (non-degenerated critical points)**

$$\square EBS^{Reg}(x_0, \dots, x_k) = \{x \in Aff(x_0, \dots, x_k), s.t. H(x, \lambda^*(x)) \neq 0\}$$

$$\square EBS^+(x_0, \dots, x_k) = \{x \in Aff(x_0, \dots, x_k), s.t. H(x, \lambda^*(x)) \text{ Pos. def.}\}$$


Theorem: EBS partitioned into cells by the index of the Hessian of  $\lambda$ -variance:  $KBS = EBS^+$  on  $M^*$

[X.P. Barycentric Subspace Analysis on Manifolds. Annals of statistics. 2018. To appear. arXiv:1607.02833]

# Example on the sphere

## Manifold

- Unit sphere  $\mathcal{M} = S_n$  embedded in  $\mathbb{R}^{n+1}$
- $\|x\| = 1$



## Exp and log map

$$\exp_x(v) = \cos(\|v\|)x + \frac{\sin(\|v\|)}{\|v\|}v$$

$$\log_x(y) = f(\theta)(y - \cos(\theta)) \quad \text{with} \quad \theta = \arccos(x^t y)$$

## Distance

$$\text{dist}(x, y) = \|\log_x(y)\| = \theta$$

## (k+1)-pointed & punctured Sphere

- $X = [x_0, x_1, \dots, x_k] \in (S_n)^k$
- Punctured sphere: exclude antipodal points:  $S_n^* = S_n / -X$

# KBS / FBS with 3 points on the sphere


**EBS: great subspheres spanned by reference points (mod cut loci)**

$$\text{EBS}(x_0, \dots, x_k) = \text{Span}(X) \cap S_n \setminus \text{Cut}(X) \quad \text{Aff}(x_0, \dots, x_k) = \text{Span}(X) \cap S_n$$

**KBS/FBS: look at index of the Hessian of  $\lambda$ -variance**

$$H(x, \lambda) = \sum \lambda_i \theta_i \cot(\theta_i) (\text{Id} - xx^t) + \sum (1 - \lambda_i \theta_i \cot(\theta_i)) \overrightarrow{xx_i} \overrightarrow{xx_i}^t$$

- Complex algebraic geometry problem [Buss & Fillmore, ACM TG 2001]
- 3 points of the n-sphere: EBS partitioned in cell complex by index of critical point
- **KBS/EBS less interesting than EBS/affine span**



Weighted Hessian index: **brown = -2 (min) = KBS** / **green = -1 (saddle)** / **blue = 0 (max)**

# Example on the hyperbolic space

## Manifold

- Unit pseudo-sphere  $\mathcal{M} = H_n$  embedded in Minkowski space  $\mathbb{R}^{1,n}$
- $\|x\|_*^2 = -x_0^2 + x_1^2 + \dots + x_n^2 = -1$


## Exp and log map

$$\exp_x(v) = \cosh(\|v\|_*)x + \frac{\sinh(\|v\|_*)}{\|v\|_*}v$$

$$\log_x(y) = f_*(\theta)(y - \cosh(\theta)) \quad \text{with} \quad \theta = \operatorname{arcosh}(-\langle x|y \rangle_*)$$

**Distance**  $\operatorname{dist}(x, y) = \|\log_x(y)\|_* = \theta$

**Punctured hyperbolic space:** no cut locus to exclude



# Example on the hyperbolic space


**EBS = Affine span: great sub-hyperboloids spanned by reference points**

$$\text{EBS}(x_0, \dots, x_k) = \text{Aff}(x_0, \dots, x_k) = \text{Span}(X) \cap H_n$$

**KBS: locus of maximal index of the Hessian of  $\lambda$ -variance**

$$H(x, \lambda) = \sum \lambda_i \theta_i \coth(J + J_{xx}^t J^t) + \sum (1 - \lambda_i \coth(\theta_i)) J \overrightarrow{xx_i} \overrightarrow{xx_i}^t J^t$$

- Complex algebraic geometry problem
- 3 points on  $H^n$ : better than for spheres, but still disconnected components



Weighted Hessian Index: **brown = -2 (min) = KBS** / **blue = 1 (saddle)**



# Geodesic subspaces are limit cases of affine span


## Theorem

- $GS(x, w_1, \dots, w_k) = \{\exp_x(\sum_i \alpha_i w_i) \text{ for } \alpha \in R^k\}$  is the limit of  $Aff(x_0, \exp_{x_0}(\epsilon w_1), \dots, \exp_{x_0}(\epsilon w_k))$  when  $\epsilon \rightarrow 0$ .
- Reference points converge to a 1<sup>st</sup> order (k,n)-jet
  - PGA [Fletcher et al. 2004, Sommer et al. 2014]
  - GPGA [Huckemann et al. 2010]

## Conjecture

- This can be generalized to higher order derivatives
  - Quadratic, cubic splines [Vialard, Singh, Niethammer]
  - Principle nested spheres [Jung, Dryden, Marron 2012]
  - Quotient of Lie group action [Huckemann, Hotz, Munk, 2010]

# Application in Cardiac motion analysis




[ Marc-Michel Rohé et al., MICCAI 2016, Media 45:1-12, 2018 ]

# Application in Cardiac motion analysis

Find weights  $\lambda_i$  and SVFs  $v_i$  such that:

- $v_i$  registers image to reference  $i$
- $\sum_i \lambda_i v_i = 0$




Optimize reference images to achieve best registration over the sequence

[ Marc-Michel Rohé et al., MICCAI 2016, Media 45:1-12, 2018 ]

# Application in Cardiac motion analysis

Optimal Reference Frames

Barycentric coefficients curves




[ Marc-Michel Rohé et al., MICCAI 2016, Media 45:1-12, 2018 ]


# Cardiac Motion Signature

Low-dimensional representation of motion using:

Optimal Reference Frames



Barycentric coefficients curves



Dimension reduction from **+10M voxels** to **3 reference frames + 60 coefficients**

Tested on **10 controls** [1] and **16 Tetralogy of Fallot** patients [2]

[1] Tobon-Gomez, C., et al.: Benchmarking framework for myocardial tracking and deformation algorithms: an open access database. *Medical Image Analysis* (2013)




[2] Mcleod K., et al.: Spatio-Temporal Tensor Decomposition of a Polyaffine Motion Model for a Better Analysis of Pathological Left Ventricular Dynamics. *IEEE TMI* (2015)

# Cardiac motion synthesis

Original Sequence

Barycentric Reconstruction  
(3 images)

PCA Reconstruction  
(2 modes)



30 images

3 images + 2 coeff.

1 image + 2 SVF + 2 coeff.

Reconstr. error: 18.75  
Compression ratio: 1/10







Reconstr. error: 26.32 (+40%)  
Compression ratio: 1/4

[ Marc-Michel Rohé et al., MICCAI 2016, Media 45:1-12, 2018 ]



# Cardiac motion tracking

Method evaluated on one **synthetic cardiac sequence\***



[\*] Prakosa, A., et al.: Generation of Synthetic but Visually Realistic Time Series of

# ***Geometric Statistics: Mathematical foundations and applications in computational anatomy***

Intrinsic Statistics on Riemannian Manifolds

Manifold-Valued Image Processing

Metric and Affine Geometric Settings for Lie Groups

Parallel Transport to Analyze Longitudinal Deformations

## **Advances Statistics: CLT & PCA**

- Estimation of the empirical Fréchet mean & CLT
- Principal component analysis in manifolds
- Natural subspaces in manifolds: barycentric subspaces
- **Rephrasing PCA with flags of subspaces**

# *The natural object for PCA: Flags of subspaces in manifolds*

## **Subspace approximations with variable dimension**

- Optimal unexplained variance  $\rightarrow$  non nested subspaces
- Nested forward / backward procedures  $\rightarrow$  not optimal
- Optimize first, decide dimension later  $\rightarrow$  Nestedness required  
**[Principal nested relations: Damon, Marron, JMIV 2014]**

## **Flags of affine spans in manifolds: $FL(x_0 < x_1 < \dots < x_n)$**

- Sequence of nested subspaces

$$Aff(x_0) \subset Aff(x_0, x_1) \subset \dots Aff(x_0, \dots x_i) \subset \dots Aff(x_0, \dots x_n) = M$$

## **Barycentric subspace analysis (BSA):**

- Energy on flags: Accumulated Unexplained Variance  
 $\rightarrow$  **optimal flags of subspaces in Euclidean spaces = PCA**

**[ X.P. Barycentric Subspace Analysis on Manifolds, Annals of Statistics 2018 ]**

# Robustness with $L_p$ norms

## Affine spans is stable to p-norms

- $\sigma^p(x, \lambda) = \frac{1}{p} \sum \lambda_i \text{dist}^p(x, x_i) / \sum \lambda_i$
- Critical points of  $\sigma^p(x, \lambda)$  are also critical points of  $\sigma^2(x, \lambda')$  with  $\lambda'_i = \lambda_i \text{dist}^{p-2}(x, x_i)$  (non-linear reparameterization of affine span)


## Unexplained p-variance of residuals

- $2 < p \rightarrow +\infty$ : more weight on the tail,  
at the limit: penalizes the maximal distance to subspace
- $0 < p < 2$ : less weight on the tail of the residual errors:  
statistically robust estimation
  - Non-convex for  $p < 1$  even in Euclidean space
  - But sample-limited algorithms do not need gradient information

# Experiments on the sphere

## 3 clusters on a 5D sphere

- 10, 9 and 8 points (stddev 6 deg) around three orthogonal axes plus 30 points uniformly samples on 5D sphere




- **FBS: Forward Barycentric Subspace:** mean and median not in clusters
- **1-PBS / 2-PBS: Pure Barycentric Subspace with backward ordering:** ok for  $k=2$  only
- **1-BSA / 2-BSA: Barycentric Subspace Analysis up to order  $k$ :** less sensitive to  $p$  &  $k$



# Experiments on the hyperbolic space

## 3 clusters on a 5D hyperboloid (50% outliers)

- 15 random points (stddev 0.015) around an equilateral triangle of length 1.57 plus 15 points of stddev 1.0 (truncated at max 1.5)



- **FBS: Forward Barycentric Subspace:** ok for  $p \leq 0.5$
- **1-PBS / 2-PBS: Pure Barycentric Subspace with backward ordering:** ok for  $k=2$  only
- **1-BSA / 2-BSA: Barycentric Subspace Analysis up to order  $k$ :** ok for  $p \leq 1$

# Take home messages

## Natural subspaces in manifolds


- PGA & Godesic subspaces:  
look at data points from the (unique) mean
- Barycentric subspaces:  
« triangulate » several reference points
  - Justification of multi-atlases?

## Critical points (affine span) rather than minima (FBS/KBS)

- Barycentric coordinates need not be positive (convexity is a problem)
- Affine notion (more general than metric)
  - Generalization to Lie groups (SVFs)?

## Natural flag structure for PCA

- Hierarchically embedded approximation subspaces to summarize / describe data





# Pushing the frontiers of Geometric Statistics

## Beyond the mean and unimodal concentrated laws


- **Flags (nested sequences) of subspace in manifolds**
- **Non Gaussian statistical models within subspaces?**

## Beyond the Riemannian / metric structure

- Riemannian manifolds, Non-Positively Curved (NPC) metric spaces
- Towards **Affine connection, Quotient, Stratified spaces**

## Unify statistical estimation theory



- Explore influence of **curvature, singularities** (borders, corners, stratifications) on non-asymptotic estimation theory



# Quotient spaces

## Functions/Images modulo time/space parameterization


- Amplitude and phase discrimination problem



[Allasoniere, Amit, Trouvé, 2005],  
Example by Loic Devillier, IPMI 2017

# Noise in top space = Bias in quotient spaces

The curvature of the **template shape's orbit** and presence of **noise** creates a repulsive bias



**Theorem [Miolane et al. (2016)]:** Bias of estimator  $\hat{T}$  of the template  $T$

$$\text{Bias}(\hat{T}, T) = \frac{\sigma^2}{2} H(T) + \mathcal{O}(\sigma^4)$$

where  $H(T)$  : **mean curvature vector of template's orbit**

Extension to Hilbert of  $\infty$ -dim: bias for  $\sigma > 0$ , asymptotic for  $\sigma \rightarrow \infty$ ,  
[Devilliers, Allasonnière, Trouvé and XP. SIIMS 2017, Entropy, 2017]

**→ Estimated atlas is topologically more complex than should be**

# Towards non-smooth spaces


## Stratified spaces

### □ Correlation matrices


- Positive semi definite (PSD) matrices with unit diagonal [Grubisic and Pietersz, 2004]

### □ Orthant spaces (phylogenetic trees)

- BHV tree space [Billera Holmes Voigt, Adv Appl Math, 2001]  
[Nye AOS 2011] [Feragen 2013] [Barden & Le, 2017]



Adapted from  
[Rousseeuw and  
Molenberghs,  
1994].






**Tree space T4**

Adapted from [Dinh et al, AoS 2018,


## Can we explain non standard statistical results?

- **Sticky mean** [Hotz et al 2013] [Barden & Le 2017], **repulsive mean** [Miolane 2017]
- **Faster convergence rate with #sample in NPC spaces** [Basrak, 2010]



[Ellingson et al, Topics in Nonparametric Statistics, 2014]

# RIEMANNIAN GEOMETRIC STATISTICS IN MEDICAL IMAGE ANALYSIS



To appear 09-2019, Elsevier

Edited by  
Xavier Pennec,  
Stefan Sommer, Tom Fletcher



## Part 1: Foundations

- 1: Riemannian geometry [Sommer, Fletcher, Pennec]
- 2: Statistics on manifolds [Fletcher]
- 3: Manifold-valued image processing with SPD matrices [Pennec]
- 4: Riemannian Geometry on Shapes and Diffeomorphisms [Marsland, Sommer]
- 5: Beyond Riemannian: the affine connection setting for transformation groups [Pennec, Lorenzi]

## Part 2: Statistics on Manifolds and Shape Spaces

- 6: Object Shape Representation via Skeletal Models (s-reps) and Statistical Analysis [Pizer, Maron]
- 7: Inductive Fréchet Mean Computation on  $S(n)$  and  $SO(n)$  with Applications [Chakraborty, Vemuri]
- 8: Statistics in stratified spaces [Ferage, Nye]
- 9: Bias in quotient space and its correction [Miolane, Devillier, Pennec]
- 10: Probabilistic Approaches to Statistics on Manifolds: Stochastic Processes, Transition Distributions, and Fiber Bundle Geometry [Sommer]
- 11: Elastic Shape Analysis, Square-Root Representations and Their Inverses [Zhang, Klassen, Srivastava]

## Part 3: Deformations, Diffeomorphisms and their Applications

- 13: Geometric RKHS models for handling curves and surfaces in Computational Anatomy : currents, varifolds, f-shapes, normal cycles [Charlie, Charon, Glaunes, Gori, Roussillon]
- 14: A Discretize-Optimize Approach for LDDMM Registration [Polzin, Niethammer, Vialad, Modezitski]
- 15: Spatially varying metrics in the LDDMM framework [Vialard, Risser]
- 16: Low-dimensional Shape Analysis In the Space of Diffeomorphisms [Zhang, Fleche, Wells, Golland]
- 17: Diffeomorphic density matching, Bauer, Modin, Joshi]

# References on Barycentric Subspace Analysis

## □ **Barycentric Subspace Analysis on Manifolds**

X. P. Annals of Statistics. 46(6A):2711-2746, 2018. [arXiv:1607.02833]

- **Barycentric Subspaces and Affine Spans in Manifolds** Geometric Science of Information GSI'2015, Oct 2015, Palaiseau, France. LNCS 9389, pp.12-21, 2015.  
Warning: change of denomination since this paper: EBS → affine span
- **Barycentric Subspaces Analysis on Spheres** Mathematical Foundations of Computational Anatomy (MFCA'15), Oct 2015, Munich, Germany. pp.71-82, 2015. <https://hal.inria.fr/hal-01203815>

## □ **Sample-limited $L_p$ Barycentric Subspace Analysis on Constant Curvature Spaces.** X.P. Geometric Sciences of Information (GSI 2017), Nov 2017, Paris, France. LNCS 10589, pp.20-28, 2017.

## □ **Low-Dimensional Representation of Cardiac Motion Using Barycentric Subspaces: a New Group-Wise Paradigm for Estimation, Analysis, and Reconstruction.** M.M Rohé, M. Sermesant and X.P. Medical Image Analysis vol 45, Elsevier, April 2018, 45, pp.1-12.

- **Barycentric subspace analysis: a new symmetric group-wise paradigm for cardiac motion tracking.** M.M Rohé, M. Sermesant and X.P. Proc of MICCAI 2016, Athens, LNCS 9902, p.300-307, Oct 2016.